



DBG=950 L1=260,L2=320 HL(M8)=2100 凯斯博 P1-2

[illegible]

序号	物件名称	抽出数量	备注
1	长抽组件	1套	
2	短抽组件	1套	
3	开关架	1件	
4	铜拉杆	7套(长1根, 第二)	带螺母及垫片



序号	代 号	名 称	数 量	材料说明	重量	比重
1	Q345E1.5	钢丝绳索具零件	1		0.03	0.03
2		开夹组件	1	铁元件	0.00	0.00
3		撑杆支架	1	角钢 $40 \times 40 \times 5$	0.32	0.32
4			2		1.02	2.04
5	GB/T 3763	限位块 40×10	2		0.00	0
6		定位销套	2	Q235A	0.04	0.00

序号	代号	名称	数量	型号	重量	备注
1	001	螺母M12	2		0.22	
2	002	左右螺母	1		5.12	
3	003	垫圈	1		0.11	
4	004	左螺母	1		0.22	
5	005	右螺母	1		0.22	
6	006	右螺母	1		0.22	
7	007	螺母	1		0.22	
8	008	螺母	1		0.22	
9	009	螺母	1		0.22	
10	010	螺母	1		0.22	
11	011	螺母	1		0.22	
12	012	螺母	1		0.22	
13	013	螺母	1		0.22	
14	014	螺母	1		0.22	
15	015	螺母	1		0.22	
16	016	螺母	1		0.22	
17	017	螺母	1		0.22	
18	018	螺母	1		0.22	
19	019	螺母	1		0.22	
20	020	螺母	1		0.22	
21	021	螺母	1		0.22	
22	022	螺母	1		0.22	
23	023	螺母	1		0.22	
24	024	螺母	1		0.22	
25	025	螺母	1		0.22	
26	026	螺母	1		0.22	
27	027	螺母	1		0.22	
28	028	螺母	1		0.22	
29	029	螺母	1		0.22	
30	030	螺母	1		0.22	
31	031	螺母	1		0.22	
32	032	螺母	1		0.22	
33	033	螺母	1		0.22	
34	034	螺母	1		0.22	
35	035	螺母	1		0.22	
36	036	螺母	1		0.22	
37	037	螺母	1		0.22	
38	038	螺母	1		0.22	
39	039	螺母	1		0.22	
40	040	螺母	1		0.22	
41	041	螺母	1		0.22	
42	042	螺母	1		0.22	
43	043	螺母	1		0.22	
44	044	螺母	1		0.22	
45	045	螺母	1		0.22	
46	046	螺母	1		0.22	
47	047	螺母	1		0.22	
48	048	螺母	1		0.22	
49	049	螺母	1		0.22	
50	050	螺母	1		0.22	
51	051	螺母	1		0.22	
52	052	螺母	1		0.22	
53	053	螺母	1		0.22	
54	054	螺母	1		0.22	
55	055	螺母	1		0.22	
56	056	螺母	1		0.22	
57	057	螺母	1		0.22	
58	058	螺母	1		0.22	
59	059	螺母	1		0.22	
60	060	螺母	1		0.22	
61	061	螺母	1		0.22	
62	062	螺母	1		0.22	
63	063	螺母	1		0.22	
64	064	螺母	1		0.22	
65	065	螺母	1		0.22	
66	066	螺母	1		0.22	
67	067	螺母	1		0.22	
68	068	螺母	1		0.22	
69	069	螺母	1		0.22	
70	070					

[illegible]

· 求半长 $a=120$ 厘米, $1=25, 2=35, 3=45, 4=55$

[illegible]

项目名称: 中和·三利·医院二期工程 12月
 号 11月 10月 9月 8月 7月 6月 5月 4月 3月 2月 1月
 11月 10月 9月 8月 7月 6月 5月 4月 3月 2月 1月

$$f_{\text{eff}}(M_p) = 2/100$$

Figure 1. Schematic representation of the experimental design. The subjects were divided into two groups: a control group and an experimental group. The control group received a standard diet, while the experimental group received a diet supplemented with 10% of the total energy from fat. The subjects were then divided into two subgroups: a control subgroup and an experimental subgroup. The control subgroup received a standard diet, while the experimental subgroup received a diet supplemented with 10% of the total energy from fat. The subjects were then divided into two subgroups: a control subgroup and an experimental subgroup. The control subgroup received a standard diet, while the experimental subgroup received a diet supplemented with 10% of the total energy from fat.

巴新+技术部: 19.5.24

